



Commutativity of the adjacency matrices of graphs

S. Akbari*, F. Moazami, A. Mohammadian

Department of Mathematical Sciences, Sharif University of Technology, P.O. Box 11365-9415, Tehran, Iran

ARTICLE INFO

Article history:

Received 1 November 2006

Accepted 2 September 2008

Available online 1 October 2008

Keywords:

Adjacency matrix

Commutativity

Graph decomposition

ABSTRACT

We say that two graphs G_1 and G_2 with the same vertex set commute if their adjacency matrices commute. In this paper, we find all integers n such that the complete bipartite graph $K_{n,n}$ is decomposable into commuting perfect matchings or commuting Hamilton cycles. We show that there are at most $n - 1$ linearly independent commuting adjacency matrices of size n ; and if this bound occurs, then there exists a Hadamard matrix of order n . Finally, we determine the centralizers of some families of graphs.

© 2008 Elsevier B.V. All rights reserved.

1. Introduction

For a graph G with the vertex set $\{v_1, \dots, v_n\}$, the *adjacency matrix* of G is the matrix $A = [a_{ij}] \in M_n(\mathbb{R})$ in which $a_{ij} = 1$, if v_i and v_j are adjacent; and otherwise, $a_{ij} = 0$. We say that two graphs G_1 and G_2 with the same vertex set *commute* if the adjacency matrices of G_1 and G_2 commute. A *graphical matrix* is a symmetric $(0, 1)$ -matrix all of whose diagonal entries are 0. If A is a graphical matrix, then the graph corresponding to A denoted by $G(A)$. The *centralizer* of a graph G , denoted $\mathcal{C}(G)$, is the set of all graphical matrices which commute with the adjacency matrix of G . The graph G is called *self-centralizer* if $G \cong G(A)$, for any non-zero matrix $A \in \mathcal{C}(G)$. For any $n \geq 2$, we use the symbols J_n and \mathbf{j}_n for the $n \times n$ matrix and the $1 \times n$ matrix all of whose entries are 1, respectively. Also the identity matrix of size n will be denoted by I_n . For any i, j , $1 \leq i, j \leq n$, we denote by E_{ij} , that element in $M_n(\mathbb{R})$ whose (i, j) -entry is 1 and whose other entries are 0. A *Hadamard matrix* of order n is a $(-1, 1)$ -matrix $H \in M_n(\mathbb{R})$ such that $HH^T = nI_n$.

For a vertex v of a graph G , the *degree* of v is the number of edges incident with v . A graph G is called *k-regular* if the degree of each vertex of G is k . A *perfect matching* M in a graph G is a 1-regular subgraph of G that contains all vertices of G . We denote the path and the cycle with n vertices by P_n and C_n , respectively. A cycle in G that contains all vertices of G is called a *Hamilton cycle*. We denote the complete graph with n vertices and the complete bipartite graph with two parts of sizes n and m , by K_n and $K_{n,m}$, respectively.

In [1], it has been proven that for any $n \geq 2$, the complete graph K_n is decomposable into commuting perfect matchings if and only if n is a power of 2. Also, it has been shown that K_n is decomposable into commuting Hamilton cycles if and only if n is a prime. In this paper, we investigate the decompositions of the complete bipartite graphs $K_{n,n}$ into commuting perfect matchings and commuting Hamilton cycles. Also, we determine the centralizers of some families of graphs.

2. Commuting decompositions of $K_{n,n}$

In the first theorem of this section, we wish to determine all integers n for which the graph $K_{n,n}$ is decomposable into commuting perfect matchings.

* Corresponding author.

E-mail addresses: s_akbari@sharif.edu (S. Akbari), f_moazami@math.sharif.ir (F. Moazami), ali_m@mehr.sharif.edu (A. Mohammadian).

Theorem 1. Let n be a natural number. The graph $K_{n,n}$ is decomposable into commuting perfect matchings if and only if n is a power of 2.

Proof. First assume that n is a power of 2. By Theorem 1 of [1], we can decompose K_n into $n-1$ commuting perfect matchings with adjacency matrices B_1, \dots, B_{n-1} . Hence $J_n - I_n = B_1 + \dots + B_{n-1}$. If A is the adjacency matrix of $K_{n,n}$, then using an appropriate labeling of the vertices of $K_{n,n}$, we may suppose that

$$A = \begin{bmatrix} 0 & J_n \\ J_n & 0 \end{bmatrix} = A_0 + \dots + A_{n-1},$$

where

$$A_0 = \begin{bmatrix} 0 & I_n \\ I_n & 0 \end{bmatrix} \quad \text{and} \quad A_i = \begin{bmatrix} 0 & B_i \\ B_i & 0 \end{bmatrix}, \quad \text{for } i = 1, \dots, n-1.$$

Clearly, for any $i = 0, \dots, n-1$, A_i is the adjacency matrix of a perfect matching of $K_{n,n}$. Moreover, since $\{B_1, \dots, B_{n-1}\}$ is a commuting set, $\{A_0, \dots, A_n\}$ is too. Therefore $K_{n,n}$ is decomposable into commuting perfect matchings. For the other direction, suppose that A_1, \dots, A_n are the adjacency matrices related to a decomposition of $K_{n,n}$ into commuting perfect matchings. With an appropriate labeling of the vertices of $K_{n,n}$, we have

$$A_i = \begin{bmatrix} 0 & B_i \\ B_i^\top & 0 \end{bmatrix} \quad \text{for } i = 1, \dots, n,$$

where $B_1 = I_n$. The equality $A_1 A_i = A_i A_1$ implies that $B_i = B_i^\top$. Moreover, for any i, j the equality $A_i A_j = A_j A_i$ holds, so we deduce that $B_i B_j = B_j B_i$. Also, since $B_1 = I_n$ and

$$\begin{bmatrix} 0 & J_n \\ J_n & 0 \end{bmatrix} = \begin{bmatrix} 0 & B_1 \\ B_1 & 0 \end{bmatrix} + \dots + \begin{bmatrix} 0 & B_n \\ B_n & 0 \end{bmatrix},$$

we conclude that $J_n - I_n = B_2 + \dots + B_n$. Since for $i = 2, \dots, n$, $G(A_i)$ is a perfect matching of $K_{n,n}$, the latter equation shows that B_i is a permutation (0, 1)-matrix all of whose diagonal entries are 0. Hence $G(B_2), \dots, G(B_n)$ are commuting perfect matchings of K_n that decompose K_n . Now, by Theorem 1 of [1], n is a power of 2. \square

Lemma 2. Let n be an odd number. If B is the adjacency matrix of the cycle C_n , then

$$A = \begin{bmatrix} 0 & B \\ B & 0 \end{bmatrix}$$

is the adjacency matrix of a Hamilton cycle in $K_{n,n}$.

Proof. Clearly, A is a graphical matrix and $G(A)$ is a 2-regular graph. Using Theorem 3.1.2 of [4], we have

$$\det(xI_{2n} - A) = \det \begin{bmatrix} xI_n & -B \\ -B & xI_n \end{bmatrix} = \det(xI_n - B) \det(xI_n + B). \quad (1)$$

Since C_n is a connected 2-regular graph, 2 is a simple eigenvalue of B . Furthermore, since n is odd, C_n is not a bipartite graph and so by Theorem 31.9 of [5], -2 is not an eigenvalue of B . Hence (1) shows that 2 is a simple eigenvalue of A . Since $G(A)$ is a 2-regular, it is a Hamilton cycle in $K_{n,n}$, as desired. \square

Let $\mathcal{D} = \{G_0, \dots, G_k\}$ be a decomposition of a graph G into some of its subgraphs. We say that \mathcal{D} is *quasi-Hamiltonian* if G_0 is a perfect matching in G and G_1, \dots, G_k are Hamilton cycles of G .

Theorem 3. Let n be an odd number. The graph $K_{n,n}$ has a commuting quasi-Hamiltonian decomposition if and only if n is a prime number.

Proof. Let $n = 2s + 1$, for some $s \geq 0$. First suppose that n is a prime number and

$$A = \begin{bmatrix} 0 & J_n \\ J_n & 0 \end{bmatrix}$$

is the adjacency matrix of $K_{n,n}$. From Corollary 2 of [1], we know that K_n is decomposable into commuting Hamilton cycles with adjacency matrices B_1, \dots, B_s . Since $J_n - I_n = B_1 + \dots + B_s$, using Lemma 2, we conclude that

$$\mathcal{P} = \left\{ G \left(\begin{bmatrix} 0 & I_n \\ I_n & 0 \end{bmatrix} \right), G \left(\begin{bmatrix} 0 & B_1 \\ B_1 & 0 \end{bmatrix} \right), \dots, G \left(\begin{bmatrix} 0 & B_s \\ B_s & 0 \end{bmatrix} \right) \right\}$$

is a quasi-Hamiltonian decomposition of $K_{n,n}$. Moreover, since $\{B_1, \dots, B_s\}$ is a commuting set, it is easily verified that every pair of graphs in \mathcal{P} commutes, as desired. For the other direction, assume that $\{A_0, \dots, A_s\}$ is a commuting quasi-Hamiltonian decomposition of $K_{n,n}$. By choosing an appropriate labeling of the vertices of $K_{n,n}$, we may suppose that

$$A_0 = \begin{bmatrix} 0 & I_n \\ I_n & 0 \end{bmatrix} \quad \text{and} \quad A_i = \begin{bmatrix} 0 & B_i \\ B_i^T & 0 \end{bmatrix}, \quad \text{for } i = 1, \dots, s.$$

Since $A_0 A_i = A_i A_0$, $B_i = B_i^T$, for any $i \geq 1$. Moreover, $J_n - I_n = B_1 + \dots + B_s$, hence B_i is a graphical matrix and obviously $G(B_i)$ is a 2-regular graph, for $i = 1, \dots, s$. Since $G(A_i)$ is a 2-regular connected graph, 2 is a simple eigenvalue of A_i . By applying (1), we conclude that the multiplicity of eigenvalue 2 of B_i is 1, for $i = 1, \dots, s$. This implies that $G(B_i)$ is a connected graph and therefore $\mathcal{Q} = \{G(B_1), \dots, G(B_s)\}$ is a decomposition of K_n into Hamilton cycles. Furthermore, $\{A_1, \dots, A_s\}$ is a commuting set, so it can be easily checked that any pair of graphs in \mathcal{Q} commutes. Now, Corollary 4 of [1] yields that n is a prime number. \square

Theorem 4. Let n be an even number. The graph $K_{n,n}$ is decomposable into commuting Hamilton cycles if and only if n is a power of 2.

Proof. For each permutation π on $\{1, \dots, m\}$, assume that $P_\pi = [p_{ij}] \in M_m(\mathbb{R})$ is the permutation matrix corresponding to π , that is, $p_{ij} = 1$ if and only if $j = \pi(i)$. Clearly, $G(P_\pi + P_\pi^T)$ is a Hamilton cycle in K_m if and only if π is a cycle of length m . First suppose that $K_{n,n}$ is decomposable into commuting Hamilton cycles. So the number of these cycles is $n/2$. Label the vertices of two parts of $K_{n,n}$ by $X = \{1, 3, \dots, 2n-1\}$ and $Y = \{2, 4, \dots, 2n\}$. With no loss of generality, we may assume that $C = G(P_\sigma + P_\sigma^T)$ is one of the Hamilton cycles appeared in the commuting decomposition of $K_{n,n}$, where $\sigma = (1 \ 2 \ \dots \ 2n)$. Let \mathcal{G} be the set of all Hamilton cycles in K_{2n} which have no common edge with C and commute with C . By Theorem 4 of [1], any element of \mathcal{G} has the form $G(P_{\sigma^k} + P_{\sigma^k}^T)$, where $k \in \{1, \dots, 2n\}$ and k and $2n$ are coprime. For any k , we have $P_{\sigma^k}^T = P_{\sigma^{2n-k}}$, so $|\mathcal{G}| \leq \phi(2n)/2$, where ϕ is Euler function. This implies that $\frac{n}{2} \leq \frac{\phi(2n)}{2}$. Let $n = 2^t n_0$, for some $t \geq 1$ and odd number n_0 . Since ϕ is an arithmetic function, $\phi(2n) = 2^t \phi(n_0)$ and thus the above inequality yields that $n_0 \leq \phi(n_0)$. From this we conclude that $n_0 = 1$, so n is a power of 2, as desired. For the other direction, suppose that n is a power of 2. Clearly,

$$\mathcal{H} = \left\{ G(P_{\sigma^k} + P_{\sigma^k}^T) \mid k \text{ is odd and } k < n \right\}$$

is a set of commuting Hamilton cycles in K_{2n} . For any k , we have $P_{\sigma^k} + P_{\sigma^k}^T = \sum_{i=1}^{2n} (E_{ii'} + E_{i'i})$, where $i' = i + k \pmod{2n}$. Then one can easily see that no pair of \mathcal{H} has common edge. Furthermore, $i \not\equiv i' \pmod{2}$ for any i , hence for each $H \in \mathcal{H}$, every edge of H meets X and Y . Since $|\mathcal{H}| = n/2$, \mathcal{H} is a decomposition of $K_{n,n}$ into commuting Hamilton cycles. \square

3. Commuting graphical matrices and Hadamard matrices

In Proposition 2.3.6 of [3], Heinze has determined the centralizers of the complete graph K_n and the complete bipartite graph $K_{n,n}$. In this section, we obtain the centralizers of some other graphs. First we find the maximum size of a linearly independent set of commuting graphical matrices.

Theorem 5. If \mathcal{A} is a linearly independent set of commuting graphical matrices in $M_n(\mathbb{R})$, then $|\mathcal{A}| \leq n-1$. Moreover, if the equality holds, then there exists a Hadamard matrix of order n . In particular, in the equality case, either $n \leq 2$ or n is divisible by 4.

Proof. Since \mathcal{A} is a commuting diagonalizable set, there exists an orthogonal matrix $P = [p_{ij}] \in M_n(\mathbb{R})$ such that $P^T \mathcal{A} P$ is a linearly independent set of diagonal matrices with trace 0. This shows that $|\mathcal{A}| \leq n-1$. Now, suppose that $|\mathcal{A}| = n-1$ and let $\mathcal{A} = \{A_1, \dots, A_{n-1}\}$. Since $\text{tr } A_j = 0$, it is easy to see that $\{I_n\} \cup P^T \mathcal{A} P$ is a linearly independent set of diagonal matrices in $M_n(\mathbb{R})$. So it is a basis for the linear space of diagonal matrices. Therefore, for $j = 1, \dots, n$, there exist real numbers $\lambda_{j0}, \dots, \lambda_{j,n-1}$ such that

$$E_{jj} = \lambda_{j0} I_n + \lambda_{j1} P^T A_1 P + \dots + \lambda_{j,n-1} P^T A_{n-1} P,$$

and so

$$P E_{jj} P^T = \lambda_{j0} I_n + \lambda_{j1} A_1 + \dots + \lambda_{j,n-1} A_{n-1}. \quad (2)$$

Since $\text{tr } A_1 = \dots = \text{tr } A_{n-1} = 0$, the relation (2) implies that $1 = \text{tr}(P E_{jj} P^T) = n \lambda_{j0}$, for $j = 1, \dots, n$. Hence $\lambda_{j0} = 1/n$, for $j = 1, \dots, n$. Furthermore, for any i, j , the (i, i) -entry of $P E_{jj} P^T$ is p_{ij}^2 . On the other hand, all diagonal entries of the right side of the equality (2) are λ_{j0} , thus for any i, j , $p_{ij}^2 = 1/n$. Now, if we put $H = n^{\frac{1}{2}} P$, then H is a $(-1, 1)$ -matrix and $H H^T = n I_n$. Therefore H is a Hadamard matrix of order n and by Theorem 18.1 of [5], either $n \leq 2$ or n is divisible by 4. \square

Remark 6. Let n be a power of 2. By Theorem 1 of [1], the complete graph K_n is decomposable into commuting perfect matchings. It is clear that the adjacency matrices of these perfect matchings are linearly independent. Hence there exists a linearly independent set of commuting graphical matrices of size $n-1$, when n is a power of 2.

4. The centralizers of some families of graphs

For any $n \geq 1$, let $\mathcal{P}_n(x)$ be the characteristic polynomial of the path P_n . In [2], Beezer has asked when a polynomial in a graphical matrix A yields a graphical matrix. In Theorem 3.1 of [2], a solution in the case that $G(A)$ is a path has been given as follows.

Theorem A. For any $n \geq 1$, let A_n be the adjacency matrix of P_n and suppose that $f(x) \in \mathbb{R}[x]$ is a non-zero polynomial of degree less than n . Then $f(A_n)$ is a graphical matrix if and only if $f(x) = \mathcal{P}_{2i-1}(x)$, for some i , $1 \leq i \leq \lfloor n/2 \rfloor$.

Theorem 7. For any $n \geq 1$, let A_n be the adjacency matrix of P_n . Then

$$\mathcal{C}(P_n) = \{\mathcal{P}_{2i-1}(A_n) | 1 \leq i \leq \lfloor n/2 \rfloor\}.$$

Proof. We know that all eigenvalues of P_n are mutually distinct. So the characteristic polynomial and the minimal polynomial of A_n are the same. Thus by Theorem 39.1.2 of [4], any matrix commuting with A_n is a polynomial in A_n . Now, by Theorem A the assertion is proved. \square

For any $n, m \geq 1$, let $W_{n,m}$ be the graph obtained from K_n by joining m new vertices to each vertex of K_n as pendant vertices. Note that the graph $W_{n,m}$ has $n(m+1)$ vertices.

Theorem 8. For any $n, m \geq 1$, the only connected graph commuting with $W_{n,m}$ is $W_{n,m}$. In addition, the graph $K_{1,m}$ is self-centralizer.

Proof. We label the vertices of $W_{n,m}$ by $1, \dots, n$ for the vertices of K_n and for any i , $1 \leq i \leq n$, we use the labels $n + (i-1)m + 1, \dots, n + im$ for the vertices that are adjacent to the i th vertex of K_n . Thus the adjacency matrix of $W_{n,m}$ is

$$A = \begin{bmatrix} J_n - I_n & I_n \otimes \mathbf{j}_m \\ I_n \otimes \mathbf{j}_m^T & 0_{nm} \end{bmatrix}.$$

Suppose that

$$B = \begin{bmatrix} X & Y \\ Y^T & Z \end{bmatrix}$$

commutes with A , where $X = [x_{ij}] \in M_n(\mathbb{R})$. The equality $AB = BA$ yields that $(I_n \otimes \mathbf{j}_m^T)Y = Y^T(I_n \otimes \mathbf{j}_m)$. It is not hard to see that the latter equality implies that there exists a symmetric $(0, 1)$ -matrix $C = [c_{ij}] \in M_n(\mathbb{R})$ such that $Y = C \otimes \mathbf{j}_m^T$. Moreover, it is easily verified that for every two matrices $R \in M_{r \times k}(\mathbb{R})$ and $S \in M_{k \times s}(\mathbb{R})$, the equality $R(S \otimes \mathbf{j}_m) = RS \otimes \mathbf{j}_m$ holds. Now, the relation $AB = BA$ yields that $(J_n - I_n)Y + (I_n \otimes \mathbf{j}_m)Z = X(I_n \otimes \mathbf{j}_m)$ and hence $(I_n \otimes \mathbf{j}_m)Z = (X - (J_n - I_n)C) \otimes \mathbf{j}_m$. Let $Z = [P_{ij}]_{1 \leq i, j \leq n}$, where P_{ij} is a $(0, 1)$ -matrix in $M_m(\mathbb{R})$. By multiplying the latter equality by $I_n \otimes \mathbf{j}_m^T$ from the right side, we have

$$[\sigma(P_{ij})] = m(X - (J_n - I_n)C), \quad (3)$$

where $\sigma(P_{ij})$ is the sum of all entries of the matrix P_{ij} . For any i , we have $\sigma(P_{ii}) = m(x_{ii} - \sum_{j \neq i} c_{ij})$. Since Z is a $(0, 1)$ -matrix and all diagonal entries of X are 0, from the latter equality we conclude that C is a diagonal matrix. Note that Z is a symmetric matrix and $\sigma(P_{ij}) = \mathbf{j}_m^T P_{ij} \mathbf{j}_m$, for any i, j . Hence by (3), we deduce that $(J_n - I_n)C$ is a symmetric matrix. This yields that $c_{11} = \dots = c_{nn}$. Obviously, if $C = 0$, then $G(B)$ is not connected. So assume that $C = I_n$. Since $\sigma(P_{ij}) \geq 0$ and X is a $(0, 1)$ -matrix, (3) implies that $X = J_n - I_n$ and thus $Z = 0$. Therefore we obtain that $A = B$, as desired. Furthermore, if $n = 1$ and $C = 0$, then from (3) we conclude that $B = 0$. This shows that $W_{1,m} = K_{1,m}$ is self-centralizer. \square

In the following, for any $n \geq 1$, F_n denotes the *friendship graph* on $2n+1$ vertices, that is, the graph formed by n triangles which have a common vertex.

Theorem 9. For any $n \geq 1$, the graph F_n is self-centralizer.

Proof. Label the vertices of F_n such that, for $i = 1, \dots, n$, $\{1, 2i, 2i+1\}$ be the vertices of the i th triangle of F_n . Let X be a non-zero graphical matrix such that $G(X)$ commutes with F_n . Suppose that Y is the adjacency matrix of F_n and

$$X = \begin{bmatrix} 0 & x \\ x^T & A \end{bmatrix},$$

where x is a $(0, 1)$ -vector in $M_{1 \times 2n}(\mathbb{R})$. Also, we have

$$Y = \begin{bmatrix} 0 & \mathbf{j}_{2n} \\ \mathbf{j}_{2n}^T & B \end{bmatrix}, \quad \text{where } B = I_n \otimes \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \in M_{2n}(\mathbb{R}).$$

Using the equality $XY = YX$, we find that $\mathbf{j}_{2n}A = xB$ and $\mathbf{j}_{2n}^T x + BA = x^T \mathbf{j}_{2n} + AB$. By multiplying the latter equality by \mathbf{j}_{2n} , we obtain that $2nx + \mathbf{j}_{2n}BA = s\mathbf{j}_{2n} + \mathbf{j}_{2n}AB$, where s is the sum of all entries of x . Using the equations $\mathbf{j}_{2n}A = xB$, $\mathbf{j}_{2n}B = \mathbf{j}_{2n}$, and $B^2 = I_n$, we have

$$2nx + xB = s\mathbf{j}_{2n} + x. \quad (4)$$

If $x = 0$, then the relation $\mathbf{j}_{2n}A = xB$ yields that $A = 0$. This is impossible, since $X \neq 0$. Thus we may assume that for some index i , the i th entry of x is 1. Moreover, for $n = 1$ the assertion is clear, so assume that $n \geq 2$. To obtain a contradiction, suppose that $s \leq 2n - 1$. The i th entry of left side of (4) is at least $2n$ and the i th entry of its right side is $s + 1$. This implies that $s \geq 2n - 1$ and so exactly one entry of x is 0. Hence there exists an index k such that the k th entry of both x and xB are 1. Now, the k th entry of the left side and the right side of (4) are $2n + 1$ and $2n$, respectively. This is impossible, so $s = 2n$. This means that $x = \mathbf{j}_{2n}$. Therefore $\mathbf{j}_{2n}A = \mathbf{j}_{2n}B = \mathbf{j}_{2n}$, hence $G(A)$ is a perfect matching with $2n$ vertices. Thus $G(X) \simeq F_n$ and the proof is complete. \square

For any $n \geq 2$ and $m \geq 3$, let $\mathcal{S}_{n,m}$ be the starlike tree which obtained by joining a vertex v to m end points of m mutually disjoint paths P_n . If we label the vertex v by 1 and all vertices with distance i of v by $\{(i-1)m+2, \dots, im+1\}$ for $i = 1, \dots, n$, then the adjacency matrix of $\mathcal{S}_{n,m}$ is

$$\mathcal{A}_{n,m} = \left[\begin{array}{c|cccc} 0 & \mathbf{j}_m & 0 & \cdots & 0 \\ \hline \mathbf{j}_m^T & 0 & I_m & & \bigcirc \\ 0 & I_m & \ddots & \ddots & \\ \vdots & & \ddots & \ddots & I_m \\ 0 & \bigcirc & & I_m & 0 \end{array} \right].$$

Theorem 10. Let $n \geq 2$ and $m \geq 3$. Then every non-zero element of $\mathcal{C}(\mathcal{S}_{n,m})$ has the form

$$\left[\begin{array}{c|cccc} 0 & \mathbf{j}_m & 0 & \cdots & 0 \\ \hline \mathbf{j}_m^T & 0 & P & & \bigcirc \\ 0 & P & \ddots & \ddots & \\ \vdots & & \ddots & \ddots & P \\ 0 & \bigcirc & & P & 0 \end{array} \right],$$

where $G(P)$ is a perfect matching on m vertices. In particular, the graph $\mathcal{S}_{n,m}$ is self-centralizer.

Proof. Let

$$X = \left[\begin{array}{c|cccc} 0 & x_1 & \cdots & x_n \\ \hline x_1^T & X_{11} & \cdots & X_{1n} \\ \vdots & \vdots & \ddots & \vdots \\ x_n^T & X_{n1} & \cdots & X_{nn} \end{array} \right]$$

be a non-zero element of $\mathcal{C}(\mathcal{S}_{n,m})$. Since the first rows of two matrices $\mathcal{A}_{n,m}X$ and $X\mathcal{A}_{n,m}$ are the same, we conclude that

$$\mathbf{j}_m X_{11} = x_2, \quad (5)$$

$$\mathbf{j}_m X_{1j} = x_{j-1} + x_{j+1}, \quad \text{for } j = 2, \dots, n-1, \quad (6)$$

and

$$\mathbf{j}_m X_{1n} = x_{n-1}. \quad (7)$$

Also, by considering the (i, j) -blocks of two sides of the equality $\mathcal{A}_{n,m}X = X\mathcal{A}_{n,m}$ for $i, j \in \{2, \dots, n+1\}$, we obtain the following equations.

$$\mathbf{j}_m^T x_1 + X_{21} = x_1^T \mathbf{j}_m + X_{12}. \quad (8)$$

$$\mathbf{j}_m^T x_j + X_{2j} = X_{1,j-1} + X_{1,j+1}, \quad \text{for } j = 2, \dots, n-1. \quad (9)$$

$$\mathbf{j}_m^T x_n + X_{2n} = X_{1,n-1}. \quad (10)$$

$$X_{i-1,j} + X_{i+1,j} = X_{i,j-1} + X_{i,j+1}, \quad \text{for any } i, j, 2 \leq i, j \leq n-1. \quad (11)$$

$$X_{i-1,n} + X_{i+1,n} = X_{i,n-1}, \quad \text{for } i = 2, \dots, n-1. \quad (12)$$

$$X_{n-1,n} = X_{n,n-1}. \quad (13)$$

For $k = 1, \dots, n$, define $\mathcal{D}_k = \{X_{s,n-k+s} \mid 1 \leq s \leq k\}$ and let $\mathcal{D}_0 = \{0_m\}$. By induction on k , we prove the following assertions for $k = 1, \dots, n-1$.

$$\mathcal{P}_k : x_{n-k+1} = 0, \mathcal{D}_{k-1} = \{0\}, \text{ and } |\mathcal{D}_{k+1}| = 1.$$

First assume that $k = 1$. By multiplying two sides of (10) by \mathbf{j}_m , we find that $mx_n + \mathbf{j}_m X_{2n} = \mathbf{j}_m X_{1,n-1}$. Using the Eq. (6) for $j = n-1$, we obtain that $(m-1)x_n + \mathbf{j}_m X_{2n} = x_{n-2}$. Since $m \geq 3$ and x_{n-2} is a $(0, 1)$ -vector, $x_n = 0$ and so by (10), \mathcal{P}_1 is proved. Now, suppose that $k \geq 2$ and \mathcal{P}_t is true for any $t, 1 \leq t \leq k-1$. If $k \geq 3$, then we first show that

$$X_{2,n-k+1} = X_{3,n-k+2} = \dots = X_{k,n-1}. \quad (14)$$

By (11), we have $X_{s-1,n-k+s} + X_{s+1,n-k+s} = X_{s,n-k+s-1} + X_{s,n-k+s+1}$, for $s = 2, \dots, k-1$. Since $|\mathcal{D}_{k-1}| = 1$, $X_{s-1,n-k+s} = X_{s,n-k+s+1}$. Thus $X_{s,n-k+s-1} = X_{s+1,n-k+s}$, for any $s, 2 \leq s \leq k-1$. This establishes (14).

Now, we show that $x_{n-k+1} = 0$. If $k \geq 3$, then since \mathcal{P}_{k-2} is true, $x_{n-k+3} = 0$. By multiplying two sides of (9) by \mathbf{j}_m for $j = n-k+1$ and using the relations (6) and (7), we find that $mx_{n-k+1} + \mathbf{j}_m X_{2,n-k+1} = \mathbf{j}_m X_{1,n-k} + \mathbf{j}_m X_{1,n-k+2} = x_{n-k-1} + x_{n-k+1} + x_{n-k+1}$, for any $k \geq 2$. By applying (14), we obtain that $(m-2)x_{n-k+1} + \mathbf{j}_m X_{k,n-1} = x_{n-k-1}$. Using the Eq. (12) for $i = k$, we have

$$X_{k,n-1} = X_{k-1,n} + X_{k+1,n}. \quad (15)$$

By $|\mathcal{D}_{k-1}| = 1$, we have $X_{k-1,n} = X_{1,n-k+2}$. Moreover, since $x_{n-k+3} = 0$, the Eq. (6) for $j = n-k+2$ implies that $\mathbf{j}_m X_{1,n-k+2} = x_{n-k+1}$. Now, by multiplying two sides of (15) by \mathbf{j}_m , we obtain that $\mathbf{j}_m X_{k,n-1} = x_{n-k+1} + \mathbf{j}_m X_{k+1,n}$. This yields that $(m-1)x_{n-k+1} + \mathbf{j}_m X_{k+1,n} = x_{n-k-1}$. Since $m \geq 3$ and x_{n-k-1} is a $(0, 1)$ -vector, the latter equation implies that $x_{n-k+1} = 0$. This yields that $X_{1,n-k+2} = 0$. From $|\mathcal{D}_{k-1}| = 1$ and $X_{1,n-k+2} \in \mathcal{D}_{k-1}$, we conclude that $\mathcal{D}_{k-1} = \{0\}$. Thus the Eq. (9) for $j = n-k+1$ implies that $X_{2,n-k+1} = X_{1,n-k}$. On the other hand, $\mathcal{D}_{k-1} = \{0\}$ yields that $X_{k-1,n} = 0$. So the Eq. (12) for $i = k$ implies that $X_{k,n-1} = X_{k+1,n}$. Now, by applying (14), we obtain that $|\mathcal{D}_{k+1}| = 1$ and so \mathcal{P}_k is true. Therefore \mathcal{P}_t holds for $t = 1, \dots, n-1$.

We have $x_2 = 0$, so by (5) we find that $X_{11} = 0$ and thus $D_n = \{0\}$. Since $\{X_{12}, X_{n-1,n}\} \subseteq D_{n-1}$, we conclude that $X_{12} = X_{n-1,n}$. Moreover, we know that $X_{n,n-1} = X_{n-1,n}^T$, therefore (13) yields that X_{12} is a symmetric matrix. Hence (8) implies that the all entries of x_1 are the same. On the other hand, using the Eq. (6) for $j = 2$, we find that $\mathbf{j}_m X_{12} = x_1 + x_3$. Since X is a non-zero matrix, $X_{12} \neq 0$. Thus the equality $x_3 = 0$ yields that $x_1 = \mathbf{j}_m$ and so every column of X_{12} has exactly one non-zero entry. But X_{12} is a symmetric matrix, hence it is a permutation matrix. Now, clearly $G(X_{12})$ is a perfect matching with m vertices and $\mathcal{S}_{n,m} \simeq G(X)$, as desired. \square

We recall that the Cartesian product $G \times H$ of two graphs G and H is the graph whose vertex set is the cartesian product of the vertex sets of G and H such that two vertices (g_1, h_1) and (g_2, h_2) are adjacent if (i) $g_1 = g_2$ and $h_1 - h_2$ is an edge of H or (ii) $h_1 = h_2$ and $g_1 - g_2$ is an edge of G . The following theorem shows that the commutativity of graphs is invariant under Cartesian product.

Theorem 11. Let $\{G_1, G_2\}$ and $\{H_1, H_2\}$ be two pairs of graphs with the same order. Then $\{G_1 \times H_1, G_2 \times H_2\}$ is a pair of commuting graphs if and only if $\{G_1, G_2\}$ and $\{H_1, H_2\}$ are commuting pairs.

Proof. Let $A_i \in M_n(\mathbb{R})$ and $B_i \in M_m(\mathbb{R})$ be the adjacency matrices of G_i and H_i , respectively, for $i = 1, 2$. Then the adjacency matrix of the graph $G_i \times H_i$ is $A_i \otimes I_m + I_n \otimes B_i$, for $i = 1, 2$. It is straightforward to see that two matrices $A_1 \otimes I_m + I_n \otimes B_1$ and $A_2 \otimes I_m + I_n \otimes B_2$ commute if and only if $(A_1 A_2 - A_2 A_1) \otimes I_m + I_n \otimes (B_1 B_2 - B_2 B_1) = 0$. Clearly, this equation is equivalent to $A_1 A_2 - A_2 A_1 = B_1 B_2 - B_2 B_1 = 0$. \square

Acknowledgments

The first and third authors are indebted to the Research Council of Sharif University of Technology for support. Also the third author wishes to express his appreciation to the Institute for Studies in Theoretical Physics and Mathematics (IPM) for support.

References

- [1] S. Akbari, A. Herman, Commuting decompositions of complete graphs, J. Combin. Des. 15 (2007) 133–142.
- [2] R.A. Beezer, On the polynomial of a path, Linear Algebra Appl. 63 (1984) 221–225.
- [3] H.A. Heinze, Applications of schur rings in algebraic combinatorics: Graphs, partial difference sets and cyclotomic schemes, Ph.D. Dissertation, University of Oldenburg, 2001.
- [4] V.V. Prasolov, Problems and Theorems in Linear Algebra, in: Translations of Mathematical Monographs, vol. 134, American Mathematical Society, Providence, RI, 1994.
- [5] J.H. van Lint, R.M. Wilson, A Course in Combinatorics, second edition, Cambridge University Press, Cambridge, 2001.